



Root research is producing valuable information about tree growth

Photo: Metla/Tapani Repo

METLA

forest · knowledge · know-how · well-being

Root research is producing valuable information about tree growth under changing climate conditions

Under changing climate conditions, new information is urgently needed concerning the functioning of roots so that we can gain a better understanding of tree growth and fluxes in the boreal forest ecosystem. This information includes details of the impact of climate change on the different tree species, forest sites, and treatments, and also on the between-year variations.

Our studies are intended to enable us to assess the root and shoot responses when roots are exposed to a variety of different stress factors at different stages in the annual cycle of trees. The root studies at Metla are conducted under both in laboratory and field conditions.

Efficient research around the year – both within and beyond the laboratory

Metla has a roots laboratory in Joensuu that, in European terms, is unique. In addition, its use is shared by Metla and the University of Eastern Finland.

The roots lab consists of specially designed growth chambers known as *dasotrons*. With these, the annual cycle of trees can be accelerated by changing the dasotron conditions, so that in a single calendar year trees can be raised for the equivalent of two growing seasons and two dormancy periods. Despite their size, it is also possible to study the responses not only of small seedlings but also of saplings.

Since the air and soil conditions can be strictly controlled, our dasotron experiments are providing us with detailed information about the functioning of roots. A variety of different methods for studying roots can now be tested experimentally prior to their application under field conditions, and the results of the laboratory studies can be validated by field studies lasting several years.

Multidisciplinary approach

Our research is concerned with the effects of low soil temperature, soil frost, drought and waterlogging on roots and mycorrhizas, and also with the ways in which changes in a root system impact on shoots. Included in our study methods are microscopy, root minirhizotron imaging (growth dynamics, longevity), high pressure flow metering of root hydraulic conductance, root electrical impedance and morphology, physiological measurements of foliage (chlorophyll fluorescence, photosynthesis, water potential, cold hardiness, trunk sap flow, electrical impedance) and biomass assessment.

Further information:

Senior Researcher Tapani Repo
Finnish Forest Research Institute, Joensuu
P.O. Box 68, FI-80101 Joensuu
email: tapani.repo@metla.fi
tel. +358 50 3913136

www.metla.fi/jo/juuristolaboratorio/juuristolaboratorio.htm



- ◆ Studies are made of the root and shoot responses of trees to different stress factors (soil frost, flood, drought).
- ◆ The annual cycle of trees can be accelerated.



Each of the four dasotrons contains four root containers. Dasotrons are chambers several meters high where the soil and air conditions can be controlled independently.

The soil temperature is controlled by glycol circulation tubes on the surface and at the bottom of the root container.



Root growth is monitored by minirhizotron imaging.

Transparent acrylic tubes are embedded in the soil. The growth and longevity of single roots can be assessed by repeated imaging in the same frames.



Minirhizotron images of roots of Norway spruce taken at the same frame at the beginning and end of the growing season. The root growth is assessed by comparing the images. The size of the images is 18 x 13.5 mm.